arXiv:1612.06499v2 [math.GR] 21 Dec 2016

Invariable generation of certain groups of piecewise linear homeomorphisms of the interval

Yoshifumi Matsuda and Shigenori Matsumoto

Abstract. Let P be the group of all the orientation preserving piecewise linear homeomorphisms of the interval $[0,1]$. Given any $a>1$, let P^a be the subgroup of P consisting of all the elements with slopes in $a^{\mathbb{Z}}$, and let $P^{\mathbb{Q}}$ be the subgroup of P consisting of all the elements with slopes and breaks in $\mathbb{Q}$. We show that the groups P, P^a, $P^{\mathbb{Q}}$, as well as Thompson group F, are invariably generated.

1. Introduction

The concept of invariable generation for a group G was introduced by J. Wiegold in [**9**].

Notation 1.1. [1] For elements g and h of a group G, and a subgroup H of G, we denote:

$$h^g = ghg^{-1}, \quad h^G = \{h^g \mid g \in G\}, \quad H^g = \{h^g \mid h \in H\}.$$

Definition 1.2. (1) A subgroup H of G is called *classful* if $H \cap g^G \neq \emptyset$ for any $g \in G$, or equivalently,

$$\bigcup_{g\in G} H^g = G. \tag{1.1}$$

(2) A group G is said to be *invariably generated* if there are no classful subgroups other than G itself.

Any finite group is invariably generated, as is shown by a counting argument on (1.1). Much easier is the fact that any abelian group is invariably generated. In [**9**], it is shown that the invariable generation is extension closed. Therefore any virtually solvable group is invariably generated. It is also projection closed. Given a prime number $p > 10^{75}$, an infinite group whose arbitrary proper nontrivial subgroup is of order p is constructed in [**8**]. Such groups are necessarily generated by arbitrary two elements not from the same proper subgroup, and is invariably

2010 *Mathematics Subject Classification.* Primary 20F65. secondary 20F05.

Key words and phrases. Invariable generation, piecewise linear homeomorphism, Thompson group.

The first author is partially supported by Grant-in-Aid for Young Scientists (B) No. 25800036, and the second by Grant-in-Aid for Scientific Research (C) No. 25400096.

[1] The convention g^h is not the same as the customary one.

generated, provided there are more than one nontrivial conjugacy classes. The Grigorchuk group [**5**] is also invariably generated [**7**].

However the invariable generation is not subgroup closed: an example is given in [**10**]. It is also not direct union closed: the group of the permutations of $\mathbb{N}$ with finite support is not invariably generated, since the stabilizer of $1 \in \mathbb{N}$ is classful. Infinite groups with one nontrivial conjugacy class, constructed in [**6**], are not invariably generated. Free groups of generators ≥ 2 are not invariably generated [**9**]. More generally, nonelementary convergence groups are not invariably generated [**3**]. Acylindrically hyperbolic groups are not invariably generated [**2**]. Invariable generation of linear groups are discussed in [**7**].

The current paper is concerned with groups of piecewise linear (PL) homeomorphisms of the interval.

THEOREM 1. *Thompson group F is invariably generated.*

Our method cannot give the finite invariance generation of F obtained in [**4**]. Denote by P the group formed by all the orientation preserving PL homeomorphisms of the unit interval $[0,1]$, and by $P^{\mathbb{Q}}$ the subgroup of P formed by elements with slopes and breaks in $\mathbb{Q}$. Fix an arbitrary real number $a > 1$. Let P^a be the subgroup of P consisting of all the elements with slopes in $a^{\mathbb{Z}}$.

THEOREM 2. *The group P^a is invariably generated.*

THEOREM 3. *The group P is invariably generated.*

THEOREM 4. *The group $P^{\mathbb{Q}}$ is invariably generated.*

The proofs of the above theorems are quite similar. In Section 2, we summarize conditions for a subgroup G of P to be invariably generated. In later sections we show that F, P^a, P and $P^{\mathbb{Q}}$ satisfy these conditions independently.

2. Conditions for invariable generation

Let G be any subgroup of the group P of all the orientation preserving PL homeomorphisms of the interval $[0,1]$. We shall raise three conditions for G to be invariably generated. Let X be a dense subset of $(0,1)$ which is left invariant by G, and let $X^* = X \cup \{0,1\}$. A closed interval $I \subset [0,1]$ is called an *X-interval* (resp. *X^*-interval*) if the endpoints of I are contained in X (resp. X^*).

DEFINITION 2.1. For an X^*-interval I, let us denote

$$G(I) = \{g|_I \mid g \in G,\ \mathrm{Supp}(g) \subset I\} \quad \text{and} \quad G|_I = \{g|_I \mid g \in G,\ g(I) = I\}.$$

The first condition is to fix the relation between G and X.

Condition A: (1) The breaks of any $g \in G$ are contained in X.
(2) The group G acts on X transitively.
(3) For any X-interval I, $G|_I = G(I)$.
(4) For any X^*-interval I, there is a PL homeomorphism $\psi_I : [0,1] \to I$ such that $\psi_I(X^*) = X^* \cap I$ and $G^{\psi_I} = G(I)$.

The other two conditions are concerned with an arbitrary classful subgroup H of G.

Condition B: Any classful subgroup H acts on X transitively.

DEFINITION 2.2. For an X^*-interval I, let us denote

$$H(I) = \{h|_I \mid h \in H,\ \mathrm{Supp}(h) \subset I\} \quad \text{and} \quad H|_I = \{h|_I \mid h \in H,\ h(I) = I\}.$$

Condition C: For any classful subgroup H, there is an X-interval I_0 such that $H|_{I_0} = G(I_0)$.

In this section, we show that if a subgroup G of P satisfies conditions A, B and C, then G is invariably generated. *Henceforth in this section, we assume G satisfies conditions A, B and C.* For $f \in G$, define $s(f) \in [0,1]$ by

$$s(f) = \sup\{s \mid f|_{[0,s]} = \mathrm{id}\}.$$

By condition A(1), $s(f)$ is contained in X^*. Let H be an arbitrary classful subgroup of G

LEMMA 2.3. *For any X^*-interval $I = [t,1] \subset [0,1]$, $H(I)$ is a classful subgroup of $G(I)$.*

PROOF: Given any $f|_I \in G(I)$ where $f \in G$ with $\mathrm{Supp}(f) \subset I$, let us show that there is $g \in G$ such that $\mathrm{Supp}(g) \subset I$ and $f^g \in H$. Since H is classful, there is $g_1 \in G$ such that $f^{g_1} \in H$. Now $s(f^{g_1}) = g_1(s(f)) \in X$. Notice that $s(f) \geq t$ since $\mathrm{Supp}(f) \subset I$. By condition B, there is $g_2 \in H$ such that $g_2(g_1(s(f))) = s(f)$. Then $f^{g_2 g_1} = (f^{g_1})^{g_2}$ is an element in H since $f^{g_1} \in H$ and $g_2 \in H$. Moreover $f^{g_2 g_1}|_I$ belongs to $H(I)$ since it satisfies $s(f^{g_2 g_1}) = s(f) \geq t$. Notice that $s(f)$ is a fixed point of $g_2 g_1$. Now by condition A(3), there is an element $g \in G$ which is the identity on $[0, s(f)]$ and is equal to $g_2 g_1$ on $[s(f), 1]$. Then we have $f^g = f^{g_2 g_1} \in H$, as is required. □

For any X^*-interval $I = [t,1]$, choose a PL homeomorphism $\psi_I : [0,1] \to I$ such that $\psi_I(X) = X \cap (t,1)$ and $G^{\psi_I} = G(I)$ (condition A(4)). Since $H(I)$ is classful in $G(I)$, $H(I)^{\psi^{-1}}$ is classful in $G(I)^{\psi^{-1}} = G$. By condition B, $H(I)^{\psi^{-1}}$ acts transitively on X. Therefore $H(I)$ acts transitively on $X \cap (t,1)$. This way we get the following lemma.

LEMMA 2.4. *The classful subgroup H acts doubly transitively on X.* □

By the same argument as Lemma 2.3, applied to the inclusion of an X-interval $I = [t,t']$ into an X^*-interval $= [t,1]$, we get the following.

LEMMA 2.5. *For any X-interval I of $(0,1)$, the group $H(I)$ is a classful subgroup of $G(I)$.* □

We shall discuss consequences of condition C. Let I_0 be an X-interval such that $H|_{I_0} = G(I_0)$.

LEMMA 2.6. *We have $H(I_0) = G(I_0)$.*

PROOF: Choose an arbitrary $f|_{I_0} \in G(I_0)$, where $f \in G$ with $\mathrm{Supp}(f) \subset I_0$. Then since $H(I_0)$ is classful in $G(I_0)$ (Lemma 2.5), there is $g \in G$ such that $\mathrm{Supp}(g) \subset I_0$ and $f^g \in H$. Since $g|_{I_0} \in G(I_0) = H|_{I_0}$, there is $h \in H$ such that $h(I_0) = I_0$ and $h|_{I_0} = g|_{I_0}$. Then since $\mathrm{Supp}(f) \subset I_0$, we have $f^g = f^h$, and hence $f = (f^g)^{h^{-1}} \in H$. This, together with the assumption $\mathrm{Supp}(f) \subset I_0$, implies that $f|_{I_0} \in H(I_0)$. □

COROLLARY 2.7. *For any X-interval I in $(0,1)$, we have $H(I) = G(I)$.*

PROOF. By double transitivity of the action of H on X (Lemma 2.4), there is $h \in H$ such that $h(I_0) = I$. Now

$$H(I) = H(I_0)^h = G(I_0)^h = G(I),$$

as is required. □

Finally we shall prove that $H = G$. Let

$$G_0 = \{g \in G \mid g'(0) = g'(1) = 1\}, \quad H_0 = \{h \in H \mid h'(0) = h'(1) = 1\}.$$

Let $\{J_n\}_{n\in\mathbb{N}}$ be an increasing sequence of X-intervals such that $\cup_n J_n = (0,1)$. We have

$$H_0 = \bigcup_{n\in\mathbb{N}} H(J_n) \text{ and } G_0 = \bigcup_{n\in\mathbb{N}} G(J_n).$$

Since by the previous lemma, $H(J_n) = G(J_n)$ for any $n \in \mathbb{N}$, we get $G_0 = H_0$.

Now for any $f \in G$, there is $g \in G$ such that $f^g \in H$. But then $f^g f^{-1} = [g, f] \in G_0 \subset H$, and therefore $f \in H$. This finishes the proof that if G satisfies conditions A, B and C, then G is invariably generated.

In the rest of the paper, we use the following terminology.

DEFINITION 2.8. For $f \in P$, an interval $[0, \epsilon]$ or $[1-\epsilon, 1]$ on which f is linear is called an *end linear zone* of f.

.

3. The group F

Let us denote by $\mathbb{Z}[2^{-1}] \subset \mathbb{R}$ the set of dyadic rationals. Thompson group F is the subgroup of P consisting of all the elements with slopes in $2^{\mathbb{Z}}$ and breaks in $\mathbb{Z}[2^{-1}]$. For F, we define X in the previous section as $X = \mathbb{Z}[2^{-1}] \cap (0,1)$. It is well known that F satisfies condition A. For A(4), we can take ψ_I to be any PL homeomorphism from $[0,1]$ to I with slopes in $2^{\mathbb{Z}}$ and breaks in $\mathbb{Z}[2^{-1}]$. See [**1**] for the existence.

DEFINITION 3.1. Define a homomorphisms $\alpha : F \to \mathbb{Z}^2$ by

$$\alpha(f) = (\log_2 f'(0), \log_2 f'(1)).$$

Notice that $f \in \mathrm{Ker}(\alpha)$ if and only if $\mathrm{Supp}(f) \subset (0,1)$. It is well known [**1**] that $\mathrm{Ker}(\alpha) = [F, F]$. Of course α is a class function: $\alpha(f) = \alpha(f^g)$. Let

$$F_{1,-1} = \{f \in F \mid f(x) > x, \forall x \in (0,1),\ \alpha(f) = (1,-1)\}.$$

Given $f \in F_{1,-1}$, points 2^{-i} from an end linear zone of f at 0 are contained in a single orbit of the $\langle f \rangle$-action. Their images by high iterates of f which lie in an end linear zone at 1 are of the form $1 - k2^{-j}$ for some positive odd integer k.

DEFINITION 3.2. Define a map $\beta : F_{1,-1} \to 2\mathbb{N} - 1$, by setting $\beta(f)$ to be the above odd integer k.

LEMMA 3.3. *The map β is surjective.*

PROOF. Let $k \in 2\mathbb{N}-1$ be given. Choose a large integer j, and define $g \in F$ by setting

$$g(x) = 2x \text{ on } [0, 2^{-j}],$$
$$g(x) = 2^{-1}(x-1)+1 \text{ on } [1-k2^{-j}.1],$$

and g is a PL homeomorphism with slopes in $2^{\mathbb{Z}}$ and breaks in $\mathbb{Z}[2^{-1}]$ from the interval $[2^{-j}, 2^{-j+1}]$ to $[2^{-j+1}, 1-k2^{-j}]$. Then we have $\beta(g) = k$. □

LEMMA 3.4. *The map β is class invariant, that is, $\beta(g^f) = \beta(g)$ for any $g \in F_{1,-1}$ and any $f \in F$.*

PROOF. Assume $\beta(g) = k \in 2\mathbb{N}-1$ for $g \in F_{1,-1}$. Then there is an orbit $\mathcal{O}$ of g which contains 2^{-j} and $1-k2^{-j}$ for any large j. Choose an arbitrary element $f \in F$ and assume that the slopes of f are 2^{j_0} near 0 and 2^{j_1} near 1. Then f maps $\mathcal{O}$ to an orbit of g^f which contains 2^{-j+j_0} and $1-k2^{-j+j_1}$ for any large j, showing that $\beta(g^f) = k$. □

Let H be an arbitrary classful subgroup of F.

COROLLARY 3.5. *The map β restricted to $H \cap F_{1,-1}$ is surjective onto $2\mathbb{N}-1$.* □

The next lemma shows that condition B of Section 2 is satisfied by F.

LEMMA 3.6. *The classful subgroup H acts transitively on $X = \mathbb{Z}[2^{-1}] \cap (0,1)$.*

PROOF. By Corollary 3.5, there is an element $h_0 \in H$ such that $\beta(h_0) = 1$. Thus for any large j, the points 2^{-j}, as well as $1-2^{-j}$, are on one orbit of h_0. Again by Corollary 3.5, the H orbit of these points contains $1-ka^{-j}$ for any $k \in 2\mathbb{N}-1$ and any large j. Applying negative iterates of h_0, we get that the H orbit contains all the points in X. □

We need more in order to establish condition C for F. For n large, let $I_n = [2^{-n-1}, 2^{-n}]$, $J_n = [1-2^{-n}, 1-2^{-n-1}]$ and let $\phi_n : [0,1] \to I_n$, $\psi_n : [0,1] \to J_n$ be the orientation preserving surjective linear map of slope 2^{-n-1}. Let

$$F_{1,-1,1} = \{g \in F_{1,-1} \mid \beta(g) = 1\}.$$

Given any $g \in F_{1,-1,1}$, if we choose n large enough, some iterate g^N maps I_n onto J_n. The map $\psi_n^{-1} \circ g^N \circ \phi_n$ is independent of the choice of n. In fact, if $k > 0$, $g^k\phi_{n+k} = \phi_n$ and $g^k\psi_n = \psi_{n+k}$. Therefore we have

$$\psi_{n+k}^{-1} g^{N+2k} \phi_{n+k} = (\psi_{n+k}^{-1} g^k) g^N (g^k \phi_{n+k}) = \psi_n^{-1} g^N \phi_n.$$

Notice also that $\psi_n^{-1} g^N \phi_n$ is an element of F.

DEFINITION 3.7. Define a map $\gamma : F_{1,-1,1} \to F$ by $\gamma(f) = \psi_n^{-1} \circ g^N \circ \phi_n$.

LEMMA 3.8. *The map γ is surjective.*

We shall adopt a bit longer proof, which is applicable also to the group P^a in the next section.

PROOF. Choose an arbitrary element $g \in F_{1,-1,1}$ which is linear on $[0, 2^{-n}]$ and $[1-2^{-n}, 1]$. There is $N > 0$ such that g^N maps I_n onto J_n. Let $f_0 = \psi_n^{-1} g^N \phi_n \in F$. Any element of F can be written as ff_0 for some $f \in F$. The map $\hat{f} = \psi_n f \psi_n^{-1}$ is a PL homeomorphism of the interval J_n with slopes in $2^{\mathbb{Z}}$ and breaks in $\mathbb{Z}[2^{-1}]$.

Define an element $g_1 \in F_{1,-1,1}$ to be equal to $\hat{f}g$ on J_{n-1} and equal to g elsewhere. Notice that $\hat{f}g$ is still linear on $[0, 2^{-n}]$ and $[1-2^{-n}, 1]$. We also have

$$\psi_n^{-1} g_1^N \phi_n = \psi_n^{-1} \hat{f} g^N \phi_n = (\psi_n^{-1} \hat{f} \psi_n)(\psi_n^{-1} g^N \phi_n) = f f_0.$$

Since ff_0 is an arbitrary element of F, we are done. □

LEMMA 3.9. *The map γ is class invariant. Precisely, if $g \in F_{1,-1,1}$ and $f \in F$, then $\gamma(g^f) = \gamma(g)$.*

PROOF. Choose n large enough so that g and f are linear on the intervals $[0, 2^{-n}]$ and $[1-2^{-n}, 1]$. Since $g \in F_{1,-1,1}$, some iterate g^N of g maps I_n to J_n. Put $k = \gamma(g) = \psi_n^{-1} g^N \phi_n$, and let us show that $\gamma(g^f) = k$. We assume f is of slope 2^{j_0} on $[0, 2^{-n}]$ and of slope 2^{j_1} on $[1-2^{-n}, 1]$. Then g^f is linear (of slope 2) on $[0, 2^{-n+j_0}]$, and is linear (of slope 2^{-1}) on $[1-2^{-n+j_1}, 1]$. The map g^f maps I_{n-j_0} onto J_{n-j_1}. Since $\phi_{n-j_0} = f\phi_n$ and $\psi_{n-j_1} = f\psi_n$, we have

$$\psi_{n-j_1}^{-1} (g^f)^N \phi_{n-j_0} = \psi_n^{-1} f^{-1} (f g f^{-1})^N f \phi_n = \psi_n^{-1} g^N \phi_n = k.$$

If n is big enough compared with j_0 and j_1, we have $\phi_{n-j_0} = (g^f)^{j_0 - j_1} \phi_{n-j_1}$. Therefore

$$k = \psi_{n-j_1}^{-1} (g^f)^N \phi_{n-j_0} = \psi_{n-j_1}^{-1} (g^f)^{N+j_0-j_1} \phi_{n-j_1}.$$

This shows $\gamma(g^f) = k$, as is required. □

COROLLARY 3.10. *The map γ restricted to $H_{1,-1,1} = H \cap F_{1.-1,1}$ is surjective onto F.* □

Fix once and for all an element $h_0 \in H_{1,-1,1}$ such that $\gamma(h_0) = \mathrm{id}$. Thus there is $n > 0$ such that h_0 is linear on $[0, 2^{-n}]$ and $[1-2^{-n}, 1]$, that some iterate h_0^N maps I_n onto J_n and that $\psi_n^{-1} h_0^N \phi_n = \mathrm{id}$. The next lemma shows that the group F satisfies condition C of Section 2.

LEMMA 3.11. *We have $H|_{I_n} = F(I_n)$.*

PROOF: Choose an arbitrary element $\hat{f} \in F(I_n)$ and let $f = \phi_n^{-1} \hat{f} \phi_n \in F$. By Corollary 3.10, there is $h_1 \in H_{1,-1,1}$ such that $\gamma(h_1) = f$. More precisely, for some big $m > 0$, there is $N > 0$ such that $h_1^N(I_m) = J_m$ and that $\psi_m^{-1} h_1^N \phi_m = f$. One can choose m to be greater than n in the lemma. Then some iterate $h_0^{N_0}$ of h_0 maps I_m onto J_m and $\psi_m^{-1} h_0^{N_0} \phi_m$ is still the identity. Thus

$$\phi_m^{-1} h_0^{-N_0} h_1^N \phi_m = (\psi_m^{-1} h_0^{N_0} \phi_m)^{-1} (\psi_m^{-1} h_1^N \phi_m) = \mathrm{id}^{-1} f = f,$$

and since $h_0^{n-m} \phi_n = \phi_m$,

$$\phi_n^{-1} h_0^{m-n} (h_0^{-N_0} h_1^N) h_0^{n-m} \phi_n = f.$$

But this means

$$h_0^{m-n} (h_0^{-N_0} h_1^N) h_0^{n-m}|_{I_n} = \hat{f}.$$

Since $\hat{f} \in F(I_n)$ is arbitrary and the LHS is in $H|_{I_n}$, we are done. □

4. The group P^a

Let $a > 1$ be an arbitrary real number.

DEFINITION 4.1. Given two compact intervals I and J, we denote by $PL^a(I, J)$ the space of the PL homeomorphisms from I to J with slopes in $a^{\mathbb{Z}}$. Such a map is called a *PL^a homeomorphism*.

LEMMA 4.2. *For any compact interval I and J, the space $PL^a(I, J)$ is nonempty.*

PROOF. Let $I = [p, q]$ and $J = [r, s]$. Consider a line $L \subset \mathbb{R}^2$ of slope a^n, $n > 1$, passing through the point (p, r), and another line L' of slope a^{-m}, $m > 1$, passing through (q, s). If n and m are sufficiently large, L and L' intersect at a point in the open rectangle $(p, q) \times (r, s)$, yielding the graph of a desired map in $PL^a(I, J)$. □

DEFINITION 4.3. Define a group P^a by $P^a = PL^a([0, 1], [0, 1])$.

We choose $X = (0, 1)$ in condition A. Then the group P^a satisfies A(1), A(2) and A(3) by virtue of Lemma 4.2. For A(4), we just take $\psi_I : [0, 1] \to I$ to be the orientation preserving linear homeomorphism. Therefore in this section, X^*-intervals are just closed intervals. In the rest we shall establish conditions B and C for P^a by almost the same method as in Section 3.

Define a homomorphism $\alpha : P^a \to \mathbb{Z}^2$ by

$$\alpha(f) = (\log_a f'(0), \log_a f'(1)).$$

Clearly α is a surjective class function. Let

$$P^a_{1,-1} = \{g \in P^a \mid g(x) > x, \forall x \in (0, 1), \alpha(g) = (1, -1)\}.$$

Given $g \in P^a_{1,-1}$, points a^{-i} for i large are contained in a single orbit of g. Consider their images by high iterates of g which are near 1. They are of the form $1 - \xi a^{-j}$ for some number $\xi \in (a^{-1}, 1]$.

DEFINITION 4.4. Define a map $\beta : P^a_{1,-1} \to (a^{-1}, 1]$, by setting $\beta(g)$ to be the above number ξ.

Then one can show that the map β is a surjective class function just as Lemmas 3.3 and 3.4 in Section 3. In particular, the map β restricted to $H \cap P^a_{1,-1}$ is surjective onto $(a^{-1}, 1]$, where H is an arbitrary classful subgroup of P^a. Then by the same method as Lemma 3.6, we get the following lemma, which establishes condition B.

LEMMA 4.5. *Any classful subgroup H acts transitively on $(0, 1)$.*

For a positive integer n, let $I_n = [a^{-n-1}, a^{-n}]$, $J_n = [1 - a^{-n}, 1 - a^{-n-1}]$ and let $\phi_n : [0, 1] \to I_n$, $\psi_n : [0, 1] \to J_n$ be the orientation preserving linear homeomorphism *of the same slope* $a^{-n}(1 - a^{-1})$. Let

$$P^a_{1,-1,1} = \{g \in P^a_{1,-1} \mid \beta(g) = 1\}.$$

Given any $g \in P^a_{1,-1,1}$, if we choose n large enough, then g is linear on the intervals $[0, a^{-n}]$ and $[1 - a^{-n}, 1]$. By the definition of $P^a_{1,-1,1}$, some iterate g^N sends I_n to J_n, and the map $\psi_n^{-1} \circ g^N \circ \phi_n$ is independent of the choice of n. Notice also that $\psi_n^{-1} g^N \phi_n$ is an element of P^a, since ϕ_n and ψ_n are linear homeomorphisms of the same slope.

DEFINITION 4.6. Define a map $\gamma : P^a_{1,-1,1} \to P^a$ by $\gamma(f) = \psi_n^{-1} \circ g^N \circ \phi_n$.

One can show that the map γ is a surjective class function just as in Lemmas 3.8 and 3.9. Fix once and for all an element $h_0 \in H \cap P^a_{1,-1,1}$ such that $\gamma(h_0) = \mathrm{id}$. Thus there is $n > 0$ such that h_0 is linear on $[0, a^{-n}]$ and $[1 - a^{-n}, 1]$, that some iterate h_0^N maps I_n onto J_n and that $\psi_n^{-1} h_0^N \phi_n = \mathrm{id}$. Just as in Lemma 3.11, we get the following lemma which establishes condition C.

LEMMA 4.7. *We have* $H|_{I_n} = P^a(I_n)$.

5. The groups P and $P^{\mathbb{Q}}$

In this section, we mainly deal with the group P of all the orientation preserving PL homeomorphisms of $[0, 1]$. In the last part, we remark one word for necessary modifications with the group $P^{\mathbb{Q}}$. For P, put $X = (0, 1)$ as in Section 4. Then condition A is trivially fulfilled. Let H be an arbitrary classful subgroup of P. First we shall establish conditions B.

LEMMA 5.1. *The group H acts transitively on $(0, 1)$.*

PROOF. There is an element $h_0 \in H$ such that $h_0'(0) = 2$ and that $h(x) > x$ for any $x \in (0, 1)$. Assume $h_0(x) = 2x$ on the interval $[0, 2^{-n}]$ for some $n > 0$. The interval $[2^{-n-1}, 2^{-n}]$ is a fundamental domain of the action of the group $\langle h_0 \rangle$. Thus it suffices to show that for any $\xi \in (2^{-1}, 1]$, there is an element of H which maps 2^{-n} to $\xi 2^{-n}$. Choose an element $h_1 \in H$ such that $h_1'(0) = \xi$. Assume h_1 is linear on an interval $[0, 2^{-m}]$ for some $m > n$. Then $h_1(2^{-m}) = \xi 2^{-m}$, and hence $h_0^{m-n} h_1 h_0^{n-m}(2^{-n}) = \xi 2^{-n}$, as is required. □

In the rest of this section, we shall establish condition C by the following lemma.

LEMMA 5.2. *For some closed interval $I_0 \subset (0, 1)$, we have $H|_{I_0} = P(I_0)$.*

For any closed interval $I \subset [0, 1]$, denote by $\phi_I : [0, 1] \to I$ the orientation preserving bijective linear map. Define

$$P_{1,-1} = \{g \in P \mid g(x) > x,\ \forall x \in (0, 1),\ g'(0) = 2,\ g'(1) = 2^{-1}\}.$$

Let I (resp. J) be a fundamental domain of $g \in P_{1,-1}$ contained in an end linear zone of g (Definition 2.8) at 0 (resp. at 1). Thus $I = [a, 2a]$ for some $a > 0$ and $J = [1 - 2b, 1 - b]$ for some $b > 0$. If there is $N > 0$ such that $g^N(I) = J$, we say that the pair I and J are *monitoring intervals for g*. The map $f = \phi_J^{-1} g^N \phi_I \in P$ is called the *information of g monitored by I and J*. We also say that *I and J monitor the information f*.

DEFINITION 5.3. For any $g \in P_{1,-1}$, denote by $\mathcal{I}(g) \subset P$ the set of all the monitored informations of g.

LEMMA 5.4. *For any $f \in P$, there is $g \in P_{1,-1}$ such that $f \in \mathcal{I}(g)$.*

PROOF. The proof is almost the same as Lemma 3.8. □

LEMMA 5.5. *Given $g \in P_{1,-1}$ and $f \in \mathcal{I}(g)$, the intervals which monitor the information f can be chosen arbitrarily near 0 and 1.*

PROOF. If the intervals I and J monitor the information f, and if $n > 0$, then clearly the intervals $g^{-n}(I)$ and $g^n(J)$ monitor the same information f. □

LEMMA 5.6. *If $g \in P_{1,-1}$ and $g_1 \in P$, then $\mathcal{I}(g^{g_1}) = \mathcal{I}(g)$.*

PROOF. Let g and g_1 be as in the lemma, and let $f \in \mathcal{I}(g)$. It suffices to show that $f \in \mathcal{I}(g^{g_1})$. By Lemma 5.5, one can choose the monitoring intervals I, J of g which monitor the information f in the end linear zones of g_1. Then $g_1(I)$ and $g_1(J)$ are monitoring intervals of g^{g_1}, with information f since $\phi_{g_1(I)} = g_1\phi_I$ and $\phi_{g_1(J)} = g_1\phi_J$. □

COROLLARY 5.7. *For any $f \in P$, there is $h \in H \cap P_{1,-1}$ such that $f \in \mathcal{I}(h)$.* □

Choose an element $h_0 \in H \cap P_{1,-1}$ so that $\mathrm{id} \in \mathcal{I}(h_0)$, and let I_0 and J_0 be monitoring intervals of h_0 with information id. That is, there is $N_0 > 0$ such that $h_0^{N_0}(I_0) = J_0$ and $\phi_{J_0}^{-1} h_0^{N_0} \phi_{I_0} = \mathrm{id}$. Let $\hat{f}$ be an arbitrary element of $P(I_0)$, and let $f = \phi_{I_0}^{-1}\hat{f}\phi_{I_0} \in P$. By Corollary 5.7, there is $h_1 \in H \cap P_{1,-1}$ such that $f \in \mathcal{I}(h_1)$. Let I_1, J_1 be the corresponding monitoring intervals: we assume $h_1^{N_1}(I_1) = J_1$ for some $N_1 > 0$ and $\phi_{J_1}^{-1} h_1^{N_1} \phi_{I_1} = f$. Put

$$I_0 = [a, 2a],\ J_0 = [1-2b, 1-b],\ I_1 = [c, 2c] \ \text{ and } \ J_1 = [1-2d, 1-d]$$

for some $a, b, c, d > 0$. Choose an element $h_2 \in H$ such that $h_2'(0) = c/a$ and $h_2'(1) = d/b$. Choose a big $n > 0$ so that both intervals $h_0^{-n}(I_0)$ and $h_0^n(J_0)$ are in the end linear zones of h_2. Direct computation shows that $h_2(h_0^{-n}(I_0)) = h_1^{-n}(I_1)$ and $h_2(h_0^n(J_0)) = h_1^n(J_1)$. See the figure.



The equality $\phi_{J_0}^{-1} h_0^{N_0} \phi_{I_0} = \mathrm{id}$ implies (cf. the proof of Lemma 5.5)

$$\phi_{h_0^n(J_0)}^{-1} h_0^{N_0+2n} \phi_{h_0^{-n}(I_0)} = \mathrm{id}.$$

Likewise we have

$$\phi_{h_1^n(J_1)}^{-1} h_0^{N_1+2n} \phi_{h_1^{-n}(I_1)} = f.$$

These equalities show

$$\phi_{h_0^{-n}(I_0)}^{-1} h_0^{-N_0-2n} \phi_{h_0^n(J_0)} \phi_{h_1^n(J_1)}^{-1} h_0^{N_1+2n} \phi_{h_1^{-n}(I_1)} = f.$$

Since h_2 is linear on the intervals $h_0^{-n}(I_0)$ and $h_0^n(J_0)$, we have $h_2\phi_{h_0^{-n}(I_0)} = \phi_{h_1^{-n}(I_1)}$ and $h_2\phi_{h_0^n(J_0)} = \phi_{h_1^n(J_1)}$. Therefore

$$\phi_{h_0^{-n}(I_0)}^{-1} h_0^{-N_0-2n} h_2^{-1} h_1^{N_1+2n} h_2 \phi_{h_0^{-n}(I_0)} = f.$$

Finally since $h_0^n \phi_{h_0^{-n}(I_0)} = \phi_{I_0}$, we get

$$\phi_{I_0}^{-1} h_0^{-N_0-n} h_2^{-1} h_1^{N_1+2n} h_2 h_0^{-n} \phi_{I_0} = f.$$

This implies

$$h_0^{-N_0-n} h_2^{-1} h_1^{N_1+2n} h_2 h_0^{-n}|_{I_0} = \hat{f}.$$

Since $\hat{f} \in P(I_0)$ is arbitrary, and the map on the LHS is from $H|_{I_0}$, the proof of Lemma 5.2 is now complete.

For the subgroup $P^{\mathbb{Q}}$ of P consisting of all the elements with slopes and breaks in $\mathbb{Q}$, we define $X = \mathbb{Q} \cap (0,1)$. The argument for $P^{\mathbb{Q}}$ is the same for P under necessary modifications.

References

[1] J. M. Belk, *Thompson's group F*, Thesis, Cornell University.

[2] M. Bestvina and K. Fujiwara, *Handlebody subgroups in a mapping class group*, preprint, arXiv:1412.7847v2.

[3] T. Gelander, *Convergence groups are not invariably generated*, Int. Math. Res. Notices **19**(2014), 9806-9814.

[4] T. Gelander, G. Golan and K. Juschenko, *Invariable generation of Thompson groups*, preprint, arXiv:1611.08264v2.

[5] R. I. Grigorchuk, *On Burnside's problem on periodic groups,* Funktional Anal. i Prilozhen **14**(1980), 53-54.

[6] G. Higman, B. H. Neumann and H. Neumenn, *Embedding theorems for groups*, J. London Math. Soc. **24**(1949), 247-254.

[7] W. M. Kantor, A. Lubotzky and A. Shalev, *Invariance generations on infinite groups*, J. Algebara **421**(2015), 296-310.

[8] A. Yu. Ol'shanskii, *Groups of bounded period with subgroups of prime order*, Algebra i Logika **21**(1982), 553-618.

[9] J. Wiegold, *Transformation groups with fixed-point-free permutations*, Arch. Math. (Basel) **27**(1976), 473-475.

[10] J. Wiegold, *Transformation groups with fixed-point-free permutations, II*, Arch. Math. (Basel) **29**(1977), 571-573.

Department of Physics and Mathematics, College of Science and Engineering, Aoyama Gakuin University, 5-10-1 Fuchinobe, Chuo-ku, Sagamihara-shi, Kanagawa, 252-5258

E-mail address: ymatsuda@gem.aoyama.ac.jp

Department of Mathematics, College of Science and Technology, Nihon University, 1-8-14 Kanda-Surugadai, Chiyoda-ku, Tokyo, 101-8308 Japan

E-mail address: matsumo@math.cst.nihon-u.ac.jp